Chapter 2

Go with the flow

> Knowing the equations and knowing the solution are two different things. Far, far away.
>
> — T.D. Lee

(R. Mainieri, P. Cvitanović and E.A. Spiegel)

We start out with a recapitulation of the basic notions of dynamics. Our aim is narrow; we keep the exposition focused on prerequisites to the applications to be developed in this text. We assume that the reader is familiar with dynamics on the level of the introductory texts mentioned in sect. 1.9, and concentrate here on developing intuition about what a dynamical system can do. It will be a coarse brush sketch–a full description of all possible behaviors of dynamical systems is beyond human ken. Anyway, for a novice there is no shortcut through this lengthy detour; a sophisticated traveler might prefer to skip this well-trodden territory and embark upon the journey at chapter 14.

fast track:
chapter 14, p. 263

2.1 Dynamical systems

In a dynamical system we observe the world as a function of time. We express our observations as numbers and record how they change with time; given sufficiently detailed information and understanding of the underlying natural laws, we see the future in the present as in a mirror. The motion of the planets against the celestial firmament provides an example. Against the daily motion of the stars from East to West, the planets distinguish themselves by moving among the fixed stars. Ancients discovered that by knowing a sequence of planet's positions–latitudes and longitudes–its future position could be predicted.

sect. 1.3

For the solar system, tracking the latitude and longitude in the celestial sphere suffices to completely specify the planet's apparent motion. All possible values for positions and velocities of the planets form the *phase space* of the system. More generally, a state of a physical system, at a given instant in time, can be represented by a single point in an abstract space called *state space* or *phase space* $\mathcal{M}$. As the system changes, so does the *representative point* in state space. We refer to the evolution of such points as *dynamics*, and the function f^t which specifies where the representative point is at time t as the *evolution rule*.

If there is a definite rule f that tells us how this representative point moves in $\mathcal{M}$, the system is said to be deterministic. For a deterministic dynamical system, the evolution rule takes one point of the state space and maps it into exactly one point. However, this is not always possible. For example, knowing the temperature today is not enough to predict the temperature tomorrow; knowing the value of a stock today will not determine its value tomorrow. The state space can be enlarged, in the hope that in a sufficiently large state space it is possible to determine an evolution rule, so we imagine that knowing the state of the atmosphere, measured over many points over the entire planet should be sufficient to determine the temperature tomorrow. Even that is not quite true, and we are less hopeful when it comes to stocks.

chapter 11

For a deterministic system almost every point has a unique future, so trajectories cannot intersect. We say 'almost' because there might exist a set of measure zero (tips of wedges, cusps, etc.) for which a trajectory is not defined. We may think such sets a nuisance, but it is quite the contrary–they will enable us to partition state space, so that the dynamics can be better understood.

Locally, the state space $\mathcal{M}$ looks like $\mathbb{R}^d$, meaning that d numbers are sufficient to determine what will happen next. Globally, it may be a more complicated manifold formed by patching together several pieces of $\mathbb{R}^d$, forming a torus, a cylinder, or some other geometric object. When we need to stress that the dimension d of $\mathcal{M}$ is greater than one, we may refer to the point $x \in \mathcal{M}$ as x_i where $i = 1, 2, 3, \ldots, d$. The evolution rule $f^t : \mathcal{M} \to \mathcal{M}$ tells us where a point x is in $\mathcal{M}$ after a time interval t.

The pair $(\mathcal{M}, f)$ constitute a *dynamical system*.

The dynamical systems we will be studying are smooth. This is expressed mathematically by saying that the evolution rule f^t can be differentiated as many times as needed. Its action on a point x is sometimes indicated by $f(x,t)$ to remind us that f is really a function of two variables: the time and a point in state space. Note that time is relative rather than absolute, so only the time interval is necessary. This follows from the fact that a point in state space completely determines all future evolution, and



Figure 2.1: A trajectory traced out by the evolution rule f^t. Starting from the state space point x, after a time t, the point is at $f^t(x)$.



Figure 2.2: The evolution rule f^t can be used to map a region $\mathcal{M}_i$ of the state space into the region $f^t(\mathcal{M}_i)$.

it is not necessary to know anything else. The time parameter can be a real variable ($t \in \mathbb{R}$), in which case the evolution is called a *flow*, or an integer ($t \in \mathbb{Z}$), in which case the evolution advances in discrete steps in time, given by *iteration* of a *map*. Actually, the evolution parameter need not be the physical time; for example, a time-stationary solution of a partial differential equation is parameterized by spatial variables. In such situations one talks of a 'spatial profile' rather than a 'flow'.

Nature provides us with innumerable dynamical systems. They manifest themselves through their trajectories: given an initial point x_0, the evolution rule traces out a sequence of points $x(t) = f^t(x_0)$, the *trajectory* through the point $x_0 = x(0)$. A trajectory is parameterized by the time t and thus belongs to $(f^t(x_0), t) \in \mathcal{M} \times \mathbb{R}$. By extension, we can also talk of the evolution of a region $\mathcal{M}_i$ of the state space: just apply f^t to every point in $\mathcal{M}_i$ to obtain a new region $f^t(\mathcal{M}_i)$, as in figure 14.2.

2.1
page 46

Because f^t is a single-valued function, any point of the trajectory can be used to label the trajectory. If we mark the trajectory by its inital point x_0, we are describing it in the *Lagrangian coordinates*. We can regard the transport of the material point at $t = 0$ to its current point $x(t) = f^t(x_0)$ as a coordinate transformation from the Lagrangian coordinates to the *Eulerian coordinates*.

The subset of points $\mathcal{M}_{x_0} \subset \mathcal{M}$ that belong to the infinite-time trajectory of a given point x_0 is called the *orbit* of x_0; we shall talk about forward orbits, backward orbits, periodic orbits, etc.. For a flow, an orbit is a smooth continuous curve; for a map, it is a sequence of points. An orbit is a *dynamically invariant* notion. While "trajectory" refers to a state $x(t)$ at time instant t, "orbit" refers to the totality of states that can be reached from x_0, with state space $\mathcal{M}$ foliated into a union of such orbits (each $\mathcal{M}_{x_0}$ labeled by a single point belonging to the set, $x_0 = x(0)$ for example).

2.1.1 A classification of possible motions?

What are the possible trajectories? This is a grand question, and there are many answers, chapters to follow offering some. Here is the first attempt

to classify all possible trajectories:

stationary:	$f^t(x) = x$	for all t
periodic:	$f^t(x) = f^{t+T_p}(x)$	for a given minimum period T_p
aperiodic:	$f^t(x) \neq f^{t'}(x)$	for all $t \neq t'$.

chapter 5

A *periodic orbit* (or a *cycle*) p is the set of points $\mathcal{M}_p \subset \mathcal{M}$ swept out by a trajectory that returns to the initial point in a finite time. Periodic orbits form a very small subset of the state space, in the same sense that rational numbers are a set of zero measure on the unit interval.

Periodic orbits and equilibrium points are the simplest examples of 'non-wandering' invariant sets preserved by dynamics. Dynamics can also preserve higher-dimensional smooth compact invariant manifolds; most commonly encountered are the M-dimensional tori of Hamiltonian dynamics, with notion of periodic motion generalized to quasiperiodic (superposition of M incommesurate frequencies) motion on a smooth torus, and families of solutions related by a continuous symmetry.

The ancients tried to make sense of all dynamics in terms of periodic motions; epicycles, integrable systems. The embarassing truth is that for a generic dynamical systems almost all motions are aperiodic. So we refine the classification by dividing aperiodic motions into two subtypes: those that wander off, and those that keep coming back.

A point $x \in \mathcal{M}$ is called a *wandering point*, if there exists an open neighborhood $\mathcal{M}_0$ of x to which the trajectory never returns

$$f^t(x) \notin \mathcal{M}_0 \qquad \text{for all} \quad t > t_{\min} \,. \tag{2.1}$$

In physics literature, the dynamics of such state is often referred to as *transient*.

Wandering points do not take part in the long-time dynamics, so your first task is to prune them from $\mathcal{M}$ as well as you can. What remains envelops the set of the long-time trajectories, or the *non-wandering set*.

For times much longer than a typical 'turnover' time, it makes sense to relax the notion of exact periodicity, and replace it by the notion of *recurrence*. A point is *recurrent* or *non-wandering* if for any open neighborhood $\mathcal{M}_0$ of x and any time t_{min} there exists a later time t, such that

$$f^t(x) \in \mathcal{M}_0 \,. \tag{2.2}$$

In other words, the trajectory of a non-wandering point reenters the neighborhood $\mathcal{M}_0$ infinitely often. We shall denote by Ω the *non–wandering set* of f, i.e., the union of all the non-wandering points of $\mathcal{M}$. The set Ω, the non–wandering set of f, is the key to understanding the long-time behavior of

a dynamical system; all calculations undertaken here will be carried out on non–wandering sets.

So much about individual trajectories. What about clouds of initial points? If there exists a connected state space volume that maps into itself under forward evolution (and you can prove that by the method of Lyapunov functionals, or several other methods available in the literature), the flow is globally contracting onto a subset of $\mathcal{M}$ which we shall refer to as the *attractor*. The attractor may be unique, or there can coexist any number of distinct attracting sets, each with its own *basin of attraction*, the set of all points that fall into the attractor under foward evolution. The attractor can be a fixed point, a periodic orbit, aperiodic, or any combination of the above. The most interesting case is that of an aperiodic recurrent attractor, to which we shall refer loosely as a *strange attractor*. We say 'loosely', as will soon become apparent that diagnosing and proving existence of a genuine, card-carrying strange attractor is a highly nontrivial undertaking.

☞ example 2.3

Conversely, if we can enclose the non–wandering set Ω by a connected state space volume $\mathcal{M}_0$ and then show that almost all points within $\mathcal{M}_0$, but not in Ω, eventually exit $\mathcal{M}_0$, we refer to the non–wandering set Ω as a *repeller*. An example of a repeller is not hard to come by–the pinball game of sect. 1.3 is a simple chaotic repeller.

It would seem, having said that the periodic points are so exceptional that almost all non-wandering points are aperiodic, that we have given up the ancients' fixation on periodic motions. Nothing could be further from truth. As longer and longer cycles approximate more and more accurately finite segments of aperiodic trajectories, we shall establish control over non–wandering sets by defining them as the closure of the union of all periodic points.

Before we can work out an example of a non–wandering set and get a better grip on what chaotic motion might look like, we need to ponder flows in a little more depth.

2.2 Flows

There is no beauty without some strangeness.
—William Blake

A *flow* is a continuous-time dynamical system. The evolution rule f^t is a family of mappings of $\mathcal{M} \to \mathcal{M}$ parameterized by $t \in \mathbb{R}$. Because t represents a time interval, any family of mappings that forms an evolution rule must satisfy:

2.2
page 46

(a) $f^0(x) = x$ (in 0 time there is no motion)

(b) $f^t(f^{t'}(x)) = f^{t+t'}(x)$ (the evolution law is the same at all times)

(c) the mapping $(x,t) \mapsto f^t(x)$ from $\mathcal{M} \times \mathbb{R}$ into $\mathcal{M}$ is continuous.

We shall often find it conveninent to represent functional composition by '$\circ$:'

☞ appendix H.1

$$f^{t+s} = f^t \circ f^s = f^t(f^s)\,. \tag{2.3}$$

The family of mappings $f^t(x)$ thus forms a continuous (forward semi-) group. Why 'semi-'group? It may fail to form a group if the dynamics is not reversible, and the rule $f^t(x)$ cannot be used to rerun the dynamics backwards in time, with negative t; with no reversibility, we cannot define the inverse $f^{-t}(f^t(x)) = f^0(x) = x$, in which case the family of mappings $f^t(x)$ does not form a group. In exceedingly many situations of interest–for times beyond the Lyapunov time, for asymptotic attractors, for dissipative partial differential equations, for systems with noise, for non-invertible maps–the dynamics cannot be run backwards in time, hence, the circumspect emphasis on *semi*groups. On the other hand, there are many settings of physical interest, where dynamics is reversible (such as finite-dimensional Hamiltonian flows), and where the family of evolution maps f^t does form a group.

For infinitesimal times, flows can be defined by differential equations. We write a trajectory as

$$x(t+\tau) = f^{t+\tau}(x_0) = f(f(x_0,t),\tau) \tag{2.4}$$

and express the time derivative of a trajectory at point $x(t)$,

$$\left.\frac{dx}{d\tau}\right|_{\tau=0} = \partial_\tau f(f(x_0,t),\tau)|_{\tau=0} = \dot{x}(t)\,. \tag{2.5}$$

2.3
page 46

as the time derivative of the evolution rule, a vector evaluated at the same point. By considering all possible trajectories, we obtain the vector $\dot{x}(t)$ at any point $x \in \mathcal{M}$. This *vector field* is a (generalized) velocity field:

$$v(x) = \dot{x}(t)\,. \tag{2.6}$$

Newton's laws, Lagrange's method, or Hamilton's method are all familiar procedures for obtaining a set of differential equations for the vector field $v(x)$ that describes the evolution of a mechanical system. Equations of mechanics may appear different in form from (2.6), as they are often involve higher time derivatives, but an equation that is second or higher order in time can always be rewritten as a set of first order equations.



Figure 2.3: (a) The two-dimensional vector field for the Duffing system (2.7), together with a short trajectory segment. (b) The flow lines. Each 'comet' represents the same time interval of a trajectory, starting at the tail and ending at the head. The longer the comet, the faster the flow in that region.

We are concerned here with a much larger world of general flows, mechanical or not, all defined by a time-independent vector field (2.6). At each point of the state space a vector indicates the local direction in which the trajectory evolves. The length of the vector $|v(x)|$ is proportional to the speed at the point x, and the direction and length of $v(x)$ changes from point to point. When the state space is a complicated manifold embedded in $\mathbb{R}^d$, one can no longer think of the vector field as being embedded in the state space. Instead, we have to imagine that each point x of state space has a different tangent plane $T\mathcal{M}_x$ attached to it. The vector field lives in the union of all these tangent planes, a space called the *tangent bundle* $\mathbf{T}\mathcal{M}$.

Example 2.1 A two-dimensional vector field $v(x)$**:** *A simple example of a flow is afforded by the unforced Duffing system*

$$\begin{aligned} \dot{x}(t) &= y(t) \\ \dot{y}(t) &= -0.15\,y(t) + x(t) - x(t)^3 \end{aligned} \tag{2.7}$$

plotted in figure 2.3. The velocity vectors are drawn superimposed over the configuration coordinates $(x(t), y(t))$ *of state space* $\mathcal{M}$*, but they belong to a different space, the* tangent bundle $\mathbf{T}\mathcal{M}$.

2.4 page 46

$$\text{If} \qquad v(x_q) = 0\,, \tag{2.8}$$

x_q is an *equilibrium point* (also referred to as a *stationary*, *fixed*, *critical*, *rest*, *stagnation* point, *zero* of the vector field v, or *steady state*), and the trajectory remains forever stuck at x_q. Otherwise the trajectory passing through x_0 at time $t = 0$ can be obtained by integrating the equations (2.6):

$$x(t) = f^t(x_0) = x_0 + \int_0^t d\tau\, v(x(\tau))\,, \qquad x(0) = x_0\,. \tag{2.9}$$



Figure 2.4: Lorenz "butterfly" strange attractor. (J. Halcrow)

We shall consider here only *autonomous* flows, i.e., flows for which the velocity field v_i is *stationary*, not explicitly dependent on time. A non-autonomous system

$$\frac{dy}{d\tau} = w(y, \tau)\,, \tag{2.10}$$

can always be converted into a system where time does not appear explicitly. To do so, extend ('suspend') state space to be $(d+1)$-dimensional by defining $x = \{y, \tau\}$, with a stationary vector field

$$v(x) = \begin{bmatrix} w(y,\tau) \\ 1 \end{bmatrix}\,. \tag{2.11}$$

2.5 page 47

The new flow $\dot{x} = v(x)$ is autonomous, and the trajectory $y(\tau)$ can be read off $x(t)$ by ignoring the last component of x.

remark 2.2

Example 2.2 Lorenz strange attractor: *Edward Lorenz arrived at the equation*

$$\dot{x} = v(x) = \begin{bmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \end{bmatrix} = \begin{bmatrix} \sigma(y-x) \\ \rho x - y - xz \\ xy - bz \end{bmatrix} \tag{2.12}$$

by a drastic simplification of the Rayleigh-Benard flow. Lorenz fixed $\sigma = 10$, $b = 8/3$, *and varied the "Rayleigh number"* ρ*. For* $0 < \rho < 1$ *the equilibrium* $EQ_0 = (0,0,0)$ *at the origin is attractive. At* $\rho = 1$ *it undergoes a pitchfork bifurcation into a pair of equilibria at*

$$x_{EQ1,2} = (\pm\sqrt{b(\rho-1)}, \pm\sqrt{b(\rho-1)}, \rho - 1)\,, \tag{2.13}$$

We shall not explore the Lorenz flow dependence on the ρ *parameter in what follows, but here is a brief synopsis: the* EQ_0 $1d$ *unstable manifold closes into a homoclinic orbit at* $\rho = 13.56\ldots$*. Beyond that, an infinity of associated periodic orbits are generated, until* $\rho = 24.74\ldots$*, where* $EQ_{1,2}$ *undergo a Hopf bifurcation.*

All computations that follow will be performed for the Lorenz parameter choice $\sigma = 10, b = 8/3, \rho = 28$*. For these parameter values the long-time dynamics is confined to the strange attractor depicted in figure 2.4. (Continued in example 3.4.)*



Figure 2.5: A trajectory of the Rössler flow at time $t = 250$. (G. Simon)

Example 2.3 The Rössler system–A flow with a strange attractor: *The Duffing flow of figure 2.3 is bit of a bore–every trajectory ends up in one of the two attractive equilibrium points. Let's construct a flow that does not die out, but exhibits a recurrent dynamics. Start with a harmonic oscillator*

$$\dot{x} = -y\,, \qquad \dot{y} = x\,. \tag{2.14}$$

The solutions are re^{it}, re^{-it}, and the whole x-y plane rotates with constant angular velocity $\dot{\theta} = 1$, period $T = 2\pi$. Now make the system unstable by adding

$$\dot{x} = -y\,, \qquad \dot{y} = x + ay\,, \qquad a > 0\,, \tag{2.15}$$

or, in radial coordinates, $\dot{r} = ar\sin^2\theta$, $\dot{\theta} = 1 + (a/2)\sin 2\theta$. The plane is still rotating with the same average angular velocity, but trajectories are now spiraling out. Any flow in the plane either escapes, falls into an attracting equilibrium point, or converges to a limit cycle. Richer dynamics requires at least one more dimension. In order to prevent the trajectory from escaping to ∞, kick it into 3rd dimension when x reaches some value c by adding

$$\dot{z} = b + z(x - c)\,, \qquad c > 0\,. \tag{2.16}$$

As x crosses c, z shoots upwards exponentially, $z \simeq e^{(x-c)t}$. In order to bring it back, start decreasing x by modifying its equation to

$$\dot{x} = -y - z\,.$$

Large z drives the trajectory toward $x = 0$; there the exponential contraction by e^{-ct} kicks in, and the trajectory drops back toward the x-y plane. This frequently studied example of an autonomous flow is called the Rössler system *(for definitiveness, we fix the parameters a, b, c in what follows):*

$$\begin{aligned} \dot{x} &= -y - z \\ \dot{y} &= x + ay \\ \dot{z} &= b + z(x - c)\,, \qquad a = b = 0.2\,, \quad c = 5.7\,. \end{aligned} \tag{2.17}$$

The system is as simple as they get–it would be linear, were it not for the sole bilinear term zx. Even for so 'simple' a system the nature of long-time solutions is far from obvious.

2.8 page 47

There are two repelling equilibrium points (2.8):

$$\begin{aligned} x_\pm &= \frac{c \pm \sqrt{c^2 - 4ab}}{2a}(a, -1, 1) \\ (x_-, y_-, z_-) &= (\,0.0070,\, -0.0351,\, 0.0351\,) \\ (x_+, y_+, z_+) &= (\,5.6929,\, -28.464,\, 28.464\,) \end{aligned} \tag{2.18}$$

One is close to the origin by construction–the other, some distance away, exists because the equilibrium condition has a 2nd-order nonlinearity.

To see what other solutions look like we need to resort to numerical integration. A typical numerically integrated long-time trajectory is sketched in figure 2.5. As we shall show in sect. 4.1, for this flow any finite volume of initial conditions shrinks with time, so the flow is contracting. *Trajectories that start out sufficiently close to the origin seem to converge to a* strange attractor. *We say 'seem' as there exists no proof that such an attractor is asymptotically aperiodic–it might well be that what we see is but a long transient on a way to an attractive periodic orbit. For now, accept that figure 2.5 and similar figures in what follows are examples of 'strange attractors.' (continued in exercise 2.8 and example 3.3)* *(Rytis Paškauskas)*

3.6 page 62

fast track: chapter 3, p. 49

2.3 Computing trajectories

> On two occasions I have been asked [by members of Parliament], 'Pray, Mr. Babbage, if you put into the machine wrong figures, will the right answers come out?' I am not able rightly to apprehend the kind of confusion of ideas that could provoke such a question.
>
> — Charles Babbage

You have not learned dynamics unless you know how to integrate numerically whatever dynamical equations you face. Sooner or later, you need to implement some finite time-step prescription for integration of the equations of motion (2.6). The simplest is the Euler integrator which advances the trajectory by $\delta\tau \times$ velocity at each time step:

$$x_i \to x_i + v_i(x)\delta\tau\,. \tag{2.19}$$

This might suffice to get you started, but as soon as you need higher numerical accuracy, you will need something better. There are many excellent reference texts and computer programs that can help you learn how to solve differential equations numerically using sophisticated numerical tools, such as pseudo-spectral methods or implicit methods. If a 'sophisticated' integration routine takes days and gobbles up terabits of memory, you are using brain-damaged high level software. Try writing a few lines of your own Runge-Kutta code in some mundane everyday language. While you

2.6 page 47

2.7 page 47

absolutely need to master the requisite numerical methods, this is neither the time nor the place to expound upon them; how you learn them is your business. And if you have developed some nice routines for solving problems in this text or can point another student to some, let us know.

2.10
page 48

Résumé

Chaotic dynamics with a low-dimensional attractor can be visualized as a succession of nearly periodic but unstable motions. In the same spirit, turbulence in spatially extended systems can be described in terms of recurrent spatiotemporal patterns. Pictorially, dynamics drives a given spatially extended system through a repertoire of unstable patterns; as we watch a turbulent system evolve, every so often we catch a glimpse of a familiar pattern. For any finite spatial resolution and finite time the system follows approximately a pattern belonging to a finite repertoire of possible patterns, and the long-term dynamics can be thought of as a walk through the space of such patterns. Recasting this image into mathematics is the subject of this book.

Commentary

Remark 2.1 Rössler and Duffing flows. The Duffing system (2.7) arises in the study of electronic circuits [2.2]. The Rössler flow (2.17) is the simplest flow which exhibits many of the key aspects of chaotic dynamics. We shall us the Rössler and the 3-pinball (see chapter 8) systems throughout ChaosBook to motivate the notions of Poincaré sections, return maps, symbolic dynamics, cyce expansions, etc., etc.. The Rössler system was introduced in ref. [2.3] as a set of equations describing no particular physical system, but capturing the essence of chaos in a simplest imaginable smooth flow. Otto Rössler, a man of classical education, was inspired in this quest by that rarely cited grandfather of chaos, Anaxagoras (456 B.C.). This, and references to earlier work can be found in refs. [2.5, 2.8, 2.11]. We recommend in particular the inimitable Abraham and Shaw illustrated classic [2.6] for its beautiful sketches of the Rössler and many other flows. Timothy Jones [2.19] has a number of interesting simulations on a Drexel website.

Remark 2.2 Lorenz equation: The Lorenz equation (2.12) is the most celebrated early illustration of "deterministic chaos" [9.17] (but not the first - the honor goes to Dame Cartwright [9.34]). Lorenz's paper, which can be found in reprint collections refs. [9.21, 9.22], is a pleasure to read, and is still one of the best introductions to the physics motivating such models. For a geophysics derivation, see Rothman course notes [2.7]. The equations, a set of ODEs in $\mathbb{R}^3$, exhibit strange attractors [9.18, 9.19, 9.20]. Frøyland [9.23] has a nice brief discussion of Lorenz flow. Frøyland and Alfsen [9.24] plot many periodic and heteroclinic orbits of the Lorenz flow; some of the symmetric ones are included in ref. [9.23]. Guckenheimer-Williams [9.25] and Afraimovich-Bykov-Shilnikov [9.26] offer in-depth discussion of the Lorenz equation. The most detailed study of the Lorenz equation was undertaken by Sparrow [9.28]. For a physical interpretation of ρ as "Rayleigh number." see Jackson [9.31] and Seydel [9.32]. Lorenz truncation to 3

modes is so drastic that the model bears no relation to the physical hydrodynamics problem that motivated it. For a detailed pictures of Lorenz invariant manifolds consult Vol II of Jackson [9.31]. Lorenz attractor is a very thin fractal – as we saw, stable manifold thinckness is of order 10^{-4} – but its fractal structure has been accurately resolved by D. Viswanath [2.9, 2.10].

(Continued in remark 9.1.)

Remark 2.3 Diagnosing chaos. In sect. 1.3.1 we have stated that a deterministic system exhibits 'chaos' if its dynamics is locally unstable (positive Lyapunov exponent) and globally mixing (positive entropy). In sect. 15.3 we shall define Lyapunov exponents, and discuss their evaluation, but already at this point it would be handy to have a few quick numerical methods to diagnose chaotic dynamics. Laskar's *frequency analysis* method [2.15] is useful for extracting quasi-periodic and weakly chaotic regions of state space in Hamiltonian dynamics with many degrees of freedom. For pointers to other numerical methods, see ref. [2.16].

Remark 2.4 Dynamical systems software: J.D. Meiss [2.13] has maintained for many years *Sci.nonlinear FAQ* which is now in part superseded by the SIAM Dynamical Systems website `www.dynamicalsystems.org`. The website glossary contains most of Meiss's FAQ plus new ones, and a up-to-date software list [2.14], with links to DSTool, xpp, AUTO, etc.. Springer on-line *Encyclopaedia of Mathematics* maintains links to dynamical systems software packages on eom.springer.de/D/d130210.htm.

References

[2.1] E.N. Lorenz, "Deterministic nonperiodic flow," *J. Atmos. Sci.* **20**, 130 (1963).

[2.2] G. Duffing, *Erzwungene Schwingungen bei veränderlicher Eigenfrequenz* (Vieweg. Braunschweig 1918).

[2.3] O. Rössler, *Phys. Lett.* **57A**, 397 (1976).

[2.4] "Rössler attractor," en.wikipedia.org/wiki/Rossler_map.

[2.5] J. Peinke, J. Parisi, O.E. Rössler, and R. Stoop, *Encounter with Chaos. Self-Organized Hierarchical Complexity in Semiconductor Experiments* (Springer, Berlin 1992).

[2.6] R.H. Abraham, C.D. Shaw, *Dynamics–The Geometry of Behavior* (Addison-Wesley, Redwood, Ca, 1992).

[2.7] D. Rothman, Nonlinear Dynamics I: Chaos (MIT OpenCourseWare 2006).

[2.8] R. Gilmore and M. Lefranc, *The Topology of Chaos* (Wiley, New York, 2002).

[2.9] D. Viswanath, "Symbolic dynamics and periodic orbits of the Lorenz attractor," *Nonlinearity* **16**, 1035 (2003).

[2.10] D. Viswanath, "The fractal property of the Lorenz attractor," *Physica D* **190**, 115 (2004).

[2.11] J.M.T. Thompson and H.B. Stewart *Nonlinear Dynamics and Chaos* (Wiley, New York, 2002).

[2.12] W.H. Press, B.P. Flannery, S.A. Teukolsky and W.T. Vetterling, *Numerical Recipes* (Cambridge University Press, 1986).

[2.13] J.D. Meiss, *Sci.nonlinear FAQ, Computational Resources*, amath.colorado.edu/faculty/jdm/faq.html.

[2.14] DSWeb Dynamical Systems Software, `www.dynamicalsystems.org`.

[2.15] J. Laskar, *Icarus* **88**, 257 (1990).

[2.16] Ch. Skokos, "Alignment indices: a new, simple method for determining the ordered or chaotic nature of orbits," *J. Phys* **A 34**, 10029 (2001).

[2.17] P. Cvitanović, "Periodic orbits as the skeleton of classical and quantum chaos," *Physica* **D 51**, 138 (1991).

[2.18] M.W. Hirsch, "The dynamical systems approach to differential equations," *Bull. Amer. Math. Soc.* **11**, 1 (1984)

[2.19] T. Jones, *Symmetry of Chaos Animations*, lagrange.physics.drexel.edu/flash.